

Correlation between Serum Albumin Level and Severity of Acute Ischemic Stroke : A Cross-sectional Study

Binod Pantha,¹ Milan Khadka,¹ Lochan Karki,¹ Parimal Koirala¹

¹Department of Medicine, National Academy of Medical Sciences, Bir Hospital, Kathmandu, Nepal.

ABSTRACT

Introduction: Stroke is a leading cause of disability with devastating consequences. Serum albumin has emerged as a significant prognostic marker in various conditions, including stroke. This study aims to investigate the relationship between serum albumin levels and stroke severity and outcomes in patients with acute ischemic stroke.

Methods: An observational cross-sectional study was conducted over six months at Bir Hospital, among patients admitted in medicine ward. Total population sampling was done. Serum albumin at the time of diagnosis was measured. Ischemic stroke severity was scored based on NIHSS score on admission and the functional outcome was evaluated as modified Rankin Scale . Finally, correlation of serum albumin with the severity and the outcome of stroke was made.

Results: The study of 37 patients showed significant correlation between serum albumin and NIHSS score. For high severity (NIHSS \geq 10), logistic regression analysis identified (OR: 0.086; 95% CI: 0.01-0.75, p-value 0.027) indicating there is strong association between serum albumin and NIHSS score (or severity). The study showed significant correlation between serum albumin and modified Rankin scale at the 0.01 level (2 tailed). The mean serum albumin level was 0.364 units higher (95% CI: 0.103 to 0.625) units higher in non-poor mRS score compared to poor mRS score. P-value of 0.003 suggested strong association between Serum albumin and modified Rankin Scale score.

Conclusion: Significant correlation of serum albumin level with clinical severity at the time of admission and with the outcome during discharge was found. Thus serum albumin has prognostic significance in acute ischemic stroke.

Keywords: acute ischemic stroke; albumin; modified rankin scale; national institute of health stroke scale.

INTRODUCTION

Stroke is a leading cause of disability, with severe and widespread impacts.¹ Serum albumin has emerged as a significant prognostic marker in various conditions, including stroke, with research indicating its association with both stroke severity and patient outcomes.^{2,3} The NIH Stroke Scale (NIHSS) is widely recognized for assessing initial stroke severity and predicting outcomes.⁴⁻⁷ The modified Rankin Scale (mRS) is used to evaluate functional disability, focusing on mobility and overall disability.⁸⁻¹⁰ Regardless of chosen cutoff point, serum albumin level remained an independent predictor of stroke outcome.^{11,12}

Studies have established a baseline NIHSS score with severity and mRankin score with the functional

outcome of stroke, which describes global disability with a focus on mobility.¹³⁻¹⁵

This study aims to investigate prognostic value of serum albumin levels in acute ischemic stroke, examining the relationship with NIHSS scores and mRS outcomes. By clarifying this, the study seeks to enhance understanding of serum albumin as a potential marker for stroke prognosis.

Correspondence: Dr. Binod Pantha, Department of Medicine, National Academy of Medical Sciences, Bir Hospital, Kathmandu Nepal, Email:binodpantha067@gmail.com, Phone: +977 9846140759.

METHODS

This is an observational cross-sectional study of six-month duration conducted at Bir Hospital in patients admitted in medicine wards. The study was done after getting ethical approval from the Institutional Review Committee (IRC) with reference number 34/20081/82. In this study, all adults aged 40 years or older presenting with history and clinical features suggestive of recent acute ischemic stroke admitted in the medicine wards in Bir Hospital within a study period were included. Total population sampling was done. Patients of age 40 years or older and both sexes with first-time ischemic stroke within 72 hours by clinical examination with ischemic lesion confirmed by CT Scan/MRI scan brain who gave informed consent from the patient or patient party were included. Patients with hemorrhagic stroke, ischemic stroke presenting after 72 hours of onset with recent severe infections, nephropathy, abnormal urinalysis, major trauma or surgery, major cardiac, hepatic, endocrinological disorders, skeletal disorders and cancer were excluded.

The National Institute of Health Stroke Scale Score (NIHSS) of all the patients at admission was calculated.¹⁶ serum albumin was sent as per the protocol. The correlation of NIHSS with admission serum albumin was calculated. The functional outcome was evaluated by a modified Rankin Scale (mRS)¹⁷ on discharge and its correlation value with admission serum albumin was calculated. At the time of discharge, the functional outcome was evaluated as . Data was entered in Microsoft Excel and descriptive analysis was performed.

RESULTS

A total of 37 patients with Ischemic Stroke were enrolled in this study. Out of 37 patients, 20 (54.05%) patients were male and 17 (45.95%) were female. (Table1).The mean age of the patients is approximately 67.74 years, and the median age is 69.5 years. The standard deviation (SD) of the ages is 11.32 years, and the interquartile range (IQR) is 10.0 years.

The average serum albumin level was 3.5± 0.40 mg/dl. In this study population with acute ischemic stroke, 30 (81.08%) had serum albumin in the range between 3-3.9 mg/dl (Table 1).

Albumin (g/dl)	Female n (%)	Male n (%)	Total n (%)
<2.5	1 (2.70)	-	1 (2.70)
2.5-2.9	1 (2.70)	-	1 (2.70)
3-3.9	12 (32.43)	18 (48.64)	30 (81.08)

4-4.4	3 (8.10)	2 (5.40)	5 (13.51)
≥ 4.5	-	-	-
Total	17 (45.95)	20 (54.05)	37 (100)

The mean NIHSS score at admission was 10.08. The NIHSS score of 21-42 at admission was observed in 1 (2.70%) of the patients with acute ischemic stroke, Similarly, 29 (78.37%) of patients had NIHSS scores between the range 5-15 (Table 2).

NIHSS	Female n (%)	Male n (%)	Total n (%)
0	0 (0)	0 (0)	0 (0)
1-4	2 (5.40)	1 (2.70)	3 (8.10)
5-15	12 (32.43)	17 (45.95)	29 (78.37)
16-20	2 (5.40)	2 (5.40)	4 (10.80)
21-42	1 (2.70)	0 (0)	1 (2.70)
Total	17 (45.95)	20 (54.05)	37 (100)

A Pearson correlation analysis revealed a moderate negative correlation between serum albumin levels and NIHSS scores (r = -0.403, p = 0.013).




Figure 1. Correlation between Serum albumin and NIHSS on admission

A Pearson correlation analysis showed a moderate to strong negative correlation between serum albumin levels and mRS scores (r = -0.530, p = 0.000).

A post-hoc power analysis was conducted to assess the statistical power of the observed correlations. For the correlation between serum albumin levels and NIHSS scores (r = -0.403), the power of the study was approximately 0.76, which is slightly below the conventionally desired power of 0.80. Similarly, for the correlation between serum albumin and the modified Rankin Scale (r = -0.530), the power of the study was

higher, at approximately 0.90.

DISCUSSION

In our study, most patients (81%) had serum albumin levels between 3-3.9 g/dl, which aligns with findings from similar research. For instance, Mahmud et al. observed that in a cohort of 50 acute ischemic stroke patients, the mean serum albumin level was 3.73 ± 0.63 g/dl.¹⁸ Their study found an inverse relationship between serum albumin levels and both stroke severity (NIHSS) and functional outcomes (mRS), indicating that lower albumin levels are linked to worse outcomes, reinforcing its prognostic role in stroke patients. Similarly, Kasundra et al. reported that 28% of patients had serum albumin between 3-3.4 g/dl, while 30% had levels between 3.5-3.9 g/dl.¹⁹

Our study reported a mean NIHSS score of 10.08 at admission, indicating moderate stroke severity. In contrast, Sandeep et al. found a mean NIHSS score of 15.85 ± 9.56 for survivors and 36.51 ± 4.19 for those who did not survive, reflecting a cohort with more severe strokes.²⁰ Notably, only 2.7% of our patients had an NIHSS score between 21-42, representing severe stroke, compared to the Cincinnati/Northern Kentucky Stroke Study, which reported a median NIHSS score of 3, indicating a prevalence of milder strokes in their cohort. These differences might be attributed to the varying patient populations and healthcare settings across the studies.²¹

In terms of functional outcomes, our study had a mean mRS score of 2.16 ± 1.21 at discharge, suggesting relatively mild to moderate disability. This contrasts with Sandeep et al., where the mean mRS score was 5.13 ± 0.3 in patients who died and 3.25 ± 1.12 in survivors. Moreover, our study's mortality rate was significantly lower (2.7%) compared to 9.6% in Sandeep et al.'s cohort, possibly due to differences in patient demographics, stroke severity, or healthcare management practices.²⁰

Kasundra et al. (2014) utilized the Barthel Index to measure functional independence one week after admission, focusing on short-term functional recovery and reported that higher serum albumin levels were associated with lower NIHSS scores at admission ($P < 0.001$) and better functional outcomes on the Barthel Index after one week ($P < 0.001$).¹⁹ Lower albumin levels correlated with higher NIHSS scores and poorer functional outcomes. Both studies confirm that lower serum albumin levels are associated with more severe stroke (higher NIHSS scores) and worse functional outcomes, supporting albumin's role as a prognostic biomarker in acute ischemic stroke.

In T. Dziedzic et al study, hypoalbuminemia, serum

albumin level of 35 g/l, was found in 321 from 705 (45.5%) which was similar result as in our study. In our study 17 (45.95%) had serum albumin of less than 3.5mg/dl.¹² Our study had smaller sample size (37 patients), Dziedzic et al. (2007) had larger sample size (705 patients) with albumin levels measured within 36 hours of stroke onset, focusing on the prevalence of hypoalbuminemia and its immediate correlations with stroke severity and inflammatory markers. They used Scandinavian Stroke Scale and various serum protein fractions to assess the severity and inflammatory status shortly after stroke onset. Both studies find that lower serum albumin levels are associated with more severe strokes. Our study links lower albumin levels with higher NIHSS scores and poorer mRS outcomes, while Dziedzic et al. link hypoalbuminemia with more severe strokes and various inflammatory markers.

Idicula et al. (2009) examined the association of serum albumin levels with outcome and mortality in ischemic stroke patients, exploring the potential neuroprotective effect of albumin among 444 patients with ischemic stroke.²¹ They found that higher serum albumin levels were independently associated with better outcomes ($OR = 1.12$, $p = 0.001$) and lower mortality ($OR = 0.88$, $p < 0.0001$) after adjusting for age, sex, and NIHSS score at admission. Both studies find that serum albumin levels are significantly associated with stroke outcomes. Our study highlights the predictive value of albumin for functional outcomes and severity at discharge, while Idicula et al. demonstrate that high albumin levels are associated with better outcomes and lower mortality over a longer follow-up period.

Sandeep et al. found that serum albumin was a significant predictor of both short-term and long-term outcomes in stroke patients, suggesting beneficial mechanisms similar to those identified in other studies.²⁰ They evaluated the prognostic significance of serum albumin in AIS, comparing it with NIHSS and mRS scores in an Observational study with a larger sample size of 135 patients, conducted at Liaquat University of Medical and Health Sciences. They reported a stronger negative correlation between serum albumin levels and both NIHSS ($r = -0.724$) and mRS scores ($r = -0.774$), with high specificity and sensitivity values in ROC curve analysis. Our study presents valuable insights with a smaller sample size, while Sandeep et al. provides a more comprehensive analysis with a larger sample, detailed statistical measures, and discussions on the mechanisms behind the observed correlations. Both studies agree on the potential of serum albumin as a useful biomarker in predicting stroke severity and outcomes, though Sandeep et al. offers more robust statistical evidence and broader implications.

Dash et al. (2020) explores the role of serum albumin as a prognostic marker in AIS, assessing its correlation with stroke severity (NIHSS) and outcomes at 1 week and 3 months (mRS) with a sample size of 100 patients (58% male, 42% female; mean age 65 ± 8.160 years).²² Our study is cross-sectional with outcome measures focused on discharge, while Dash et al. (2020) is prospective with extended follow-up, potentially allowing for the observation of trends over time. Both studies underscore the prognostic significance of serum albumin in AIS, with lower levels associated with worse outcomes. Both use NIHSS and mRS as primary measures of stroke severity and functional outcomes, yet the extended follow-up and larger cohort in Dash et al. (2020) offer additional insights into the long-term predictive value of serum albumin.

Nair et al. conducted a prospective observational study with 101 patients, focusing on those admitted within seven days of AIS. Stroke severity was measured using the NIHSS at admission, and functional outcomes were assessed using the Barthel index on the seventh day and third month post-stroke.²³ They identified hypoalbuminemia in 18.8% of patients. They employed unpaired t-tests to compare serum albumin levels between different dependency groups as defined by the Barthel index, both on the seventh day and third month after stroke. They found that serum albumin levels were significantly higher in patients with moderate or slight dependency compared to those with total or severe dependency at the third month post-stroke ($p = 0.008$). Similar to our study, they concluded that lower serum albumin levels were associated with greater functional dependency at three months post-stroke, though the association was not significant at the seventh day.

A large-scale cohort study using data from the Third China National Stroke Registry (CNSR-III) by Zhou et al. (2020) included 13,618 patients with acute ischemic stroke (AIS) or transient ischemic attack (TIA) and utilized multiple logistic regression models and Cox regression models to assess the association between serum albumin levels and clinical outcomes.¹¹ They concluded that low serum albumin levels are independently predictive of poor functional outcomes and mortality in patients with AIS and TIA. This provided robust evidence due to the large sample size and long-term follow-up, reinforcing the role of serum albumin as a critical prognostic factor.

A retrospective observational study involving 105 patients hospitalized after an acute stroke event by Kisialiou et al. (2022) examined a broader range of biomarkers, including ESR, fibrinogen, platelets, triglycerides, and albumin while investigating the association between these biomarkers and the size and location of ischemic lesions they found that higher albumin levels were correlated with smaller infarct sizes, suggesting a protective role, while other biomarkers were associated with different clinical outcomes.²⁴ Both studies found that higher serum albumin levels are associated with better outcomes in stroke patients, suggesting its potential as a protective factor and prognostic biomarker in acute ischemic stroke. Our study supports the existing evidence, indicating that serum albumin is an independent predictor of functional outcomes in acute ischemic stroke. The consistency of our results with previous studies reinforces the need for further research into the underlying mechanisms through which serum albumin influences stroke outcomes.

One limitation of this study is the sample size of 37 patients, which may be slightly smaller than ideal for detecting moderate correlations, especially between serum albumin levels and NIHSS scores. Based on our sample size calculation, a larger cohort, particularly for the NIHSS correlation, may have provided more robust results. This limitation suggests that the study's power to fully detect moderate correlations could be impacted, and as such, the findings should be interpreted with this consideration in mind. Despite this, the significant results observed in this study are consistent with previously published research, indicating that serum albumin levels may play a significant role in predicting stroke severity and outcomes. Future studies with larger sample sizes are recommended to confirm these findings and strengthen the statistical power. Since ischemic stroke is a condition with long-term effects and the functional outcome changes with time, lack of follow up data may weaken clinical applicability. So future studies with longitudinal study design with larger sample size should be done.

CONCLUSION

A moderate negative correlation was observed between serum albumin and NIHSS score while strong to moderate negative correlation was observed between serum albumin and mRS score.

Acknowledgement: The authors would like to thank the medical and nursing staff at Bir Hospital for their support and assistance in collecting data for this study. We are also grateful to the patients and their families for their participation in this research. Special thanks to the Department of medicine including neuromedicine for providing guidance and feedback throughout the course of the study.

Conflict of interest: The author declared no competing interests.

Author's contribution: All authors contributed equally to the study.

Ethical Consideration: Ethical Issues (Including Plagiarism, Data Fabrication, Double Publication) have been completely observed by authors.

Funding: None

REFERENCES

1. Matsumoto N, Whisnant JP, Kurland LT, Okazaki H. Natural history of stroke in Rochester, Minnesota, 1955 through 1969: an extension of a previous study, 1945 through 1954. *Stroke*. 1973;4(1):20-9. [[PubMed](#) | [Full Text](#) | [DOI](#)]
2. Gao J, Zhao Y, Du M, Guo H, Wan T, Wu M, Liu L, Wang H, Yin Q, Liu X. Serum albumin levels and clinical outcomes among ischemic stroke patients treated with endovascular thrombectomy. *Neurodegener Dis Treat*. 2021;17:401-411. [[PubMed](#) | [Full Text](#) | [DOI](#)]
3. Zhou H, Wang A, Meng X, Lin J, Jiang Y, Jing J, Zuo Y, Wang Y, Zhao X, Li H, Wang Y. Low serum albumin levels predict poor outcome in patients with acute ischemic stroke or transient ischemic attack. *Stroke Vasc Neurol*. 2021;6. [[PubMed](#) | [Full Text](#) | [DOI](#)]
4. Meyer BC, Lyden PD. The Modified National Institutes of Health Stroke Scale: Its Time Has Come. *Int J Stroke*. 2009;4(4):267-273. [[PubMed](#) | [Full Text](#) | [DOI](#)]
5. Goldstein LB, Samsa GP. Reliability of the National Institutes of Health Stroke Scale: Extension to non-neurologists in the context of a clinical trial. *Stroke*. 1997 Feb;28(2):307-10. [[PubMed](#) | [Full Text](#) | [DOI](#)]
6. Adams HP Jr, Davis PH, Leira EC, Chang KC, Bendixen BH, Clarke WR, Woolson RF, Hansen MD. Baseline NIH Stroke Scale score strongly predicts outcome after stroke: A report of the Trial of Org 10172 in Acute Stroke Treatment (TOAST). *Neurology*. 1999 Sep 22;53(1):126-31. [[PubMed](#) | [Full Text](#) | [DOI](#)]
7. Schlegel D, Kolb SJ, Luciano JM, Tovar JM, Cucchiara BL, Liebeskind DS, Kasner SE. Utility of the NIH Stroke Scale as a predictor of hospital disposition. *Stroke*. 2003 Jan;34(1):134-7. [[PubMed](#) | [Full Text](#) | [DOI](#)]
8. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke*. 1988 Nov;19(5):604-7. doi: 10.1161/01.str.19.5.604. [[PubMed](#) | [Full Text](#) | [DOI](#)]
9. Saver JL, Filip B, Hamilton S, Yanes A, Craig S, Cho M, Starkman S. Improving the reliability of stroke disability grading in clinical trials and clinical practice: the Rankin Focused Assessment (RFA). *Stroke*. 2010 May;41(5):992-5. [[PubMed](#) | [Full Text](#) | [DOI](#)]
10. Wilson JT, Hareendran A, Grant M, Baird T, Schulz UG, Muir KW, Bone I. Improving the assessment of outcomes in stroke: use of a structured interview to assign grades on the modified Rankin Scale. *Stroke*. 2002 Sep;33(9):2243-6. [[PubMed](#) | [Full Text](#) | [DOI](#)]
11. Zhou H, Wang A, Meng X, et al. Low serum albumin levels predict poor outcome in patients with acute ischaemic stroke or transient ischaemic attack. *Stroke Vasc Neurol*. 2021;6. [[PubMed](#) | [Full Text](#) | [DOI](#)]
12. Dzedzic T, Slowik A, Szczudlik A. Serum albumin level as a predictor of ischemic stroke outcome. *Stroke*. 2004 Jun;35(6):e156-8. [[PubMed](#) | [Full Text](#) | [DOI](#)]
13. Broderick JP, Adeoye O, Elm J. "Evolution of the Modified Rankin Scale and Its Use in Future Stroke Trials." *Stroke*. 2017 Jul;48(7):2007-2012. [[PubMed](#) | [Full Text](#) | [DOI](#)]
14. Lyden P, Lu M, Jackson C, et al. "Underlying Structure of the National Institutes of Health Stroke Scale: Results of a Factor Analysis." *Stroke*. 1999 Nov;30(11):2347-2354. [[PubMed](#) | [Full Text](#) | [DOI](#)]
15. Maskey A, Parajuli M, Kohli S. A study of risk factors of stroke in patients admitted to Manipal Teaching Hospital, Pokhara. *Kathmandu Univ Med J (KUMJ)*. 2012;9(4):244-7. [[PubMed](#) | [Full Text](#) | [DOI](#)]
16. Broderick JP, Adeoye O, Elm J. Evolution of the modified Rankin scale and its use in future stroke trials. *Stroke*. 2017 Jul;48(7):2007-12. [[PubMed](#) | [Full Text](#) | [DOI](#)]
17. Lyden P, Lu M, Jackson C, Marler J, Kothari R, Brott T, et al. Underlying structure of the National Institutes of Health Stroke Scale: Results of a factor analysis. *Stroke*. 1999 Nov;30(11):2347-54. [[Full Text](#) | [DOI](#)]
18. El-Din Mahmud MS, EL-Khawas HM, El Din Mahmoud Afify H, Tawfik MM. Correlation of serum albumin level with short term outcome of acute ischemic stroke patients. *QJM*. 2021 Oct 1;114(Supplement_1):hcab102.019. [[Full Text](#)]
19. Kasundra G, Sood I. Prognostic Significance Of Serum

- Albumin Levels In Acute Ischemic Stroke: Prognostic Significance Of Serum Albumin. *Natl J Integr Res Med* [Internet]. 2014 Apr. 30 [[Full Text](#)]
20. Sandeep F, Bai P, Kumari R, Shiwani S, Kumar S, Poonam F. Prognostic significance of serum albumin in acute ischemic stroke. *Eur J Pharm Med Res*. 2017;4(8):138-144. [[Full Text](#)]
 21. Idicula TT, Waje-Andreassen U, Brogger J, Naess H, Thomassen L. Serum albumin in ischemic stroke patients: the higher the better. *The Bergen Stroke Study*. *Cerebrovasc Dis*. 2009;28(1):13-7. [[PubMed](#) | [Full Text](#) | [DOI](#)]
 22. Dash PK, Behera S, Sahoo NC, Rattan R, Tripathy SK. Serum albumin level as prognostic indicator of acute ischemic stroke in tertiary care hospital admitted patients. *Int J Adv Med*. [[Full Text](#) | [DOI](#)]
 23. Nair R, Radhakrishnan K, Chatterjee A, Gorthi SP, Prabhu VA. Serum albumin as a predictor of functional outcomes following acute ischemic stroke. *J Vasc Interv Neurol*. 2018 Nov;10(2):65-68. [[PubMed](#) | [Full Text](#) | [DOI](#)]
 24. Kisialiou A, Pelone G, Carrizzo A, Grillea G, Trimarco V, Marino M, et al. Blood biomarkers role in acute ischemic stroke patients: higher is worse or better? *Immun Ageing*. 2012 Oct 31;9(1):22. [[PubMed](#) | [Full Text](#) | [DOI](#)]

© The Author(s) 2024.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>